


PRODUCT INFORMATION

IL-1 alpha (Interleukin-1 alpha), Mouse

Catalog number	C02001-5UG / C02001-20UG / C02001-100UG
Package	5 µg / 20 µg / 100 µg
Description	Interleukin 1 alpha (IL-1α) also known as hematopoietin 1 is a cytokine of the interleukin 1 family that in humans is encoded by the IL1A gene. IL-1α is produced mainly by activated macrophages, as well as neutrophils, epithelial cells, and endothelial cells. It possesses metabolic, physiological, haematopoietic activities, and plays one of the central roles in the regulation of the immune responses. It binds to the interleukin-1 receptor. It is on the pathway that activates tumor necrosis factor-alpha.
Source	<i>Escherichia coli</i>
Sequence	MSAPYTYQSDLRKLMKLVQRQKFMNDSLNTIYQDVDPKHYLSTTWLNDLQQE VKFDMYAYSSGGDDSKYPVTLKISDSQLFVSAQGEDQPVLLKELPETPKLITGS ETDLIFFWKSINSKNYFTSAAYPELFATKEQSRVHLARGLPMSMTDFQIS with polyhistidine tag at the C-terminus
Endotoxin level	<0.1 EU per 1 µg of the protein by the LAL method.
Activity	Measure by its ability to induce D10.G4.1 cells proliferation. The ED ₅₀ for this effect is <5 pg/mL. The specific activity of recombinant mouse IL-1 alpha is > 2 x 10 ⁸ IU/mg.
Purity	>95% as determined by SDS-PAGE. Ni-NTA chromatography
Formulation	The protein was lyophilized from a solution containing 1X PBS, pH 7.4.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 100 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
Storage	Lyophilized protein should be stored at -20°C. Upon reconstitution, protein aliquots should be stored at -20°C or -80°C.
Note	Please use within one month after protein reconstitution.



SDS-PAGE analysis of recombinant mouse IL-1 alpha

For Research Use Only.