


PRODUCT INFORMATION  
**IL-24 (Interleukin-24), Mouse**

<b>Catalog number</b>	C02028-5UG / C02028-20UG / C02028-100UG
<b>Package</b>	5 µg / 20 µg / 100 µg
<b>Description</b>	<p>Interleukin-24 (IL-24) is a cytokine belonging to the IL-10 family of cytokines that signals through two heterodimeric receptors: IL-20R1/IL-20R2 and IL-22R1/IL-20R2. This interleukin is also known as melanoma differentiation-associated 7 (mda-7) due to its discovery as a tumour suppressing protein. IL-24 appears to control in cell survival and proliferation by inducing rapid activation of particular transcription factors called STAT1 and STAT3. This cytokine is predominantly released by activated monocytes, macrophages and T helper 2 (Th2) cells and acts on non-haematopoietic tissues such as skin, lung and reproductive tissues. IL-24 performs important roles in wound healing, arthritis, psoriasis and cancer. Several studies have shown that cell death occurs in cancer cells/cell lines following exposure to IL-24. The gene for IL-24 is located on chromosome 1 in humans.</p>
<b>Source</b>	<i>Escherichia coli</i>
<b>Sequence</b>	<p>MQEFRFGSCQVTGVVLPPELWEAFWTVKNTVQTQDDITSIRLLKPQVLRNVSGA          ESCYLAHSLKLFYLN TVFKNYHSKIAKFKVLR SFSTLANNFIVIMSQLQPSKDNS          MLPIS ESAHQRFLLFRRAFKQLDTEVALVKAFGEVDILLTWMQK FYHL with          polyhistidine tag at the C-terminus</p>
<b>Endotoxin level</b>	<0.1 EU per 1 µg of the protein by the LAL method.
<b>Activity</b>	Measure by its ability to induce proliferation in BaF3 cells transfected with human IL-20 R alpha and human IL-20 R beta. The ED <sub>50</sub> for this effect is <0.3 ng/mL.
<b>Purity</b>	>95% as determined by SDS-PAGE. Ni-NTA chromatography
<b>Formulation</b>	The protein was lyophilized from a solution containing 1X PBS, pH 7.4.
<b>Reconstitution</b>	It is recommended to reconstitute the lyophilized protein in sterile H <sub>2</sub> O to a concentration not less than 100 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
<b>Storage</b>	Lyophilized protein should be stored at -20°C. Upon reconstitution, protein aliquots should be stored at -20°C or -80°C.
<b>Note</b>	Please use within one month after protein reconstitution.



SDS-PAGE analysis of recombinant mouse IL-24

For Research Use Only.