


BMP-15 (Bone morphogenetic protein-15), Human

v. 231001

Catalog number	C01076-5UG / C01076-20UG / C01076-100UG
Package	5 µg / 20 µg / 100 µg
Description	Bone morphogenetic protein 15 is a protein, that in humans, is encoded by the BMP15 gene. It's mainly involved in folliculogenesis. The protein encoded by this gene is a member of the TGF-β superfamily. It is a paracrine signaling molecule involved in oocyte and follicular development. Using Northern blot analysis, BMP15 has been shown to be exclusively expressed in the ovaries. This protein may be involved in oocyte maturation and follicular development as a homodimer or by forming heterodimers with a related protein, Gdf9.
Source	<i>Escherichia coli</i>
Sequence	MQADGISAEVTASSSKHSGPENNQCSLHPFQISFRQLGWDHWIAPPFYTPNY CKGTCLRVLRLDGLNSPNHAIQNLINQLVDQSVPRPSCVPYKYVPISVLMIEANG SILYKEYEGMIAESCTCR with polyhistidine tag at the C-terminus
Endotoxin level	<0.1 EU per 1 µg of the protein by the LAL method.
Activity	Measure by its ability to induce alkaline phosphatase production by ATDC5 cells. The ED ₅₀ for this effect is <17 ng/mL.
Purity	>98% as determined by SDS-PAGE.
Form	Lyophilized
Storage Buffer	Lyophilized from a 0.2 µm filtered solution containing 20 mM sodium citrate and 0.2 M NaCl, pH 3.5.
Reconstitution	It is recommended to reconstitute the lyophilized protein in 4 mM HCl to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
Stability & Storage	This product is stable after storage at: <ul style="list-style-type: none"> -20°C for 12 months in lyophilized state from date of receipt. -20°C or -80°C for 1 month under sterile conditions after reconstitution. Avoid repeated freeze/thaw cycles.



SDS-PAGE analysis of recombinant human BMP-15

For research use only.