


FGF-20 (Fibroblast growth factor-20), Human

v. 231001

| Catalog number | C01110-5UG / C01110-20UG / C01110-100UG |
|---------------------|---|
| Package | 5 µg / 20 µg / 100 µg |
| Description | FGF-20 is a member of the fibroblast growth factor (FGF) family. FGF family members possess broad mitogenic and cell survival activities, and are involved in a variety of biological processes, including embryonic development, cell growth, morphogenesis, tissue repair, tumor growth and invasion. FGF-20 signals through FGFR 2c and 3c, and is expressed during limb and brain development. Recombinant Human FGF-20 is a 23.2 kDa protein containing 209 amino acid residues. |
| Source | Escherichia coli |
| Sequence | MPLAEVGGFLGGLEGLGQQVGSHFLLPPAGERPPLLGERRSAAERSARGGPG AAQLAHLHGILRRRQLYCRTGFHLQILPDGSVQGTRQDHSLFGILEFISVAVGLV SIRGVDSGLYLGMNDKGELYGSEKLTSECIFREQFEENWYNTYSSNIYKHGDT GRRYFVALNKDGTPRDGARSKRHQKFTHFLPRPVDPERVPELYKDLLMYT with polyhistidine tag at the C-terminus |
| Endotoxin level | <0.1 EU per 1 μ g of the protein by the LAL method. |
| Activity | Measure by its ability to induce 3T3 cells proliferation. The ED ₅₀ for this effect is 1.3-3.2 ng/mL. The specific activity of recombinant human FGF-20 is > 2 x 10^5 IU/mg. |
| Purity | >98% as determined by SDS-PAGE. |
| Form | Lyophilized |
| Storage Buffer | Lyophilized from a 0.2 μ m filtered solution of PBS, pH 8.0. |
| Reconstitution | It is recommended to reconstitute the lyophilized protein in sterile H_2O to a concentration not less than 200 μ g/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved. |
| Stability & Storage | This product is stable after storage at: -20°C for 12 months in lyophilized state from date of receipt. -20°C or -80°C for 1 month under sterile conditions after reconstitution. Avoid repeated freeze/thaw cycles. |
| | |





For research use only.