


BDNF (Brain-derived neurotrophic factor), Human

v. 231001

Catalog number	C01148-5UG / C01148-20UG / C01148-100UG
Package	5 µg / 20 µg / 100 µg
Description	BDNF, also known as Brain-derived neurotrophic factor, is encoded by the BDNF Gene in human. BDNF is a member of the neurotrophin family of growth factors, which are related to the canonical nerve growth factor. Neurotrophic factors are found in the brain and the periphery. The versatility of BDNF is emphasized by its contribution to a range of adaptive neuronal responses including long-term potentiation (LTP), long-term depression (LTD), certain forms of short-term synaptic plasticity, as well as homeostatic regulation of intrinsic neuronal excitability.
Source	<i>Escherichia coli</i>
Sequence	MHSDPARRGELSVCDSEWVTAADKKTAVDMSGGTVTVLEKVPVSKGQLKQ YFYETKCNPMGYTKEGCRGIDKRHWNSQCRTTQSYVRALTMDSKKRIGWRFI RIDTSCVCTLTIKRGR with polyhistidine tag at the C-terminus
Endotoxin level	<0.1 EU per 1 µg of the protein by the LAL method.
Activity	Measure by its ability to induce proliferation in BaF3 cells transfected with TrkB. The ED ₅₀ for this effect is <2 ng/mL.
Purity	>98% as determined by SDS-PAGE.
Form	Lyophilized
Storage Buffer	Lyophilized from a 0.2 µm filtered solution containing 20 mM sodium citrate and 0.2 M NaCl, pH 3.5.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
Stability & Storage	This product is stable after storage at: <ul style="list-style-type: none"> -20°C for 12 months in lyophilized state from date of receipt. -20°C or -80°C for 1 month under sterile conditions after reconstitution. Avoid repeated freeze/thaw cycles.



SDS-PAGE analysis of recombinant human BDNF

For research use only.