


**BMP-9 (Bone morphogenetic protein-9), Human**

v. 231101

<b>Catalog number</b>	C01070-5UG / C01070-20UG / C01070-100UG
<b>Package</b>	5 µg / 20 µg / 100 µg
<b>Description</b>	BMP-9 is a member of the BMP subgroup of the TGF-beta superfamily proteins that signal through heterodimeric complexes composed of type I and type II BMP receptors. BMP-9 regulates the development and function of a variety of embryonal and adult tissues
<b>Source</b>	<i>Escherichia coli</i>
<b>Sequence</b>	SAGAGSHCQKTSLRVNFEDIGWDSWIIAPKEYEAYECKGGCFFPLADDVTPTK HAIVQTLVHLKFPTKVGKACCVPTKLSPISVLYKDDMGVPTLKYHYEGMSVAEC GCR with polyhistidine tag at the N-terminus
<b>Endotoxin level</b>	<0.01 EU per 1 µg of the protein by the LAL method.
<b>Activity</b>	Measure by its ability to induce alkaline phosphatase production by ATDC5 cells. The ED <sub>50</sub> for this effect is <0.4 ng/mL.
<b>Purity</b>	>98% as determined by SDS-PAGE.
<b>Form</b>	Lyophilized
<b>Storage Buffer</b>	Lyophilized from a 0.2 µm filtered solution containing 20 mM sodium citrate and 0.2 M NaCl, pH 3.5.
<b>Reconstitution</b>	It is recommended to reconstitute the lyophilized protein in 4 mM HCl to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
<b>Stability &amp; Storage</b>	This product is stable after storage at: <ul style="list-style-type: none"> <li>-20°C for 12 months in lyophilized state from date of receipt.</li> <li>-20°C or -80°C for 1 month under sterile conditions after reconstitution.</li> </ul> Avoid repeated freeze/thaw cycles.



SDS-PAGE analysis of recombinant human BMP-9

*For research use only.*