


IL-1 alpha (Interleukin-1 alpha), Swine

v. 231101

Catalog number	C03001-5UG / C03001-20UG / C03001-100UG
Package	5 µg / 20 µg / 100 µg
Description	Interleukin 1 alpha (IL-1α) also known as hematopoietin 1 is a cytokine of the interleukin 1 family that in humans is encoded by the IL1A gene. IL-1α is produced mainly by activated macrophages, as well as neutrophils, epithelial cells, and endothelial cells. It possesses metabolic, physiological, haematopoietic activities, and plays one of the central roles in the regulation of the immune responses. It binds to the interleukin-1 receptor. It is on the pathway that activates tumor necrosis factor-alpha.
Source	<i>Escherichia coli</i>
Sequence	MSATYSFQSNMKYNFMRVINHQICILNDARNQSIIRDPSGQYLMAAVLNNLDEAV KFDMAAYTSNDDSQLPVTLRISETRLFVSAQNEDEPVLLKELPETPKTIKDETSL LFFWEKHGNMDYFKSAAHPKLFIAATRQEKLHVHMAPGLPSVTDVFQILENQS with polyhistidine tag at the C-terminus
Endotoxin level	<0.1 EU per 1 µg of the protein by the LAL method.
Activity	Measure by its ability to induce D10.G4.1 cells proliferation. The ED ₅₀ for this effect is <50 pg/mL.
Purity	>98% as determined by SDS-PAGE.
Form	Lyophilized
Storage Buffer	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
Stability & Storage	This product is stable after storage at: <ul style="list-style-type: none"> • -20°C for 12 months in lyophilized state from date of receipt. • -20°C or -80°C for 1 month under sterile conditions after reconstitution. Avoid repeated freeze/thaw cycles.



SDS-PAGE analysis of recombinant swine IL-1 alpha

For research use only.