


Catalog number	C03007-5UG / C03007-20UG / C03007-100UG
Package	5 µg / 20 µg / 100 µg
Description	Interleukin 10 (IL-10), also known as human cytokine synthesis inhibitory factor (CSIF), is an anti-inflammatory cytokine. In humans, interleukin 10 is encoded by the IL10 gene. IL-10 signals through a receptor complex consisting of two IL-10 receptor-1 and two IL-10 receptor-2 proteins. Consequently, the functional receptor consists of four IL-10 receptor molecules. IL-10 binding induces STAT3 signalling via the phosphorylation of the cytoplasmic tails of IL-10 receptor 1 + IL-10 receptor 2 by JAK1 and Tyk2 respectively.
Source	<i>Escherichia coli</i>
Sequence	MSIKSENSCIHFPTSLPHMLRELRAAFGPVKSFFQTKDQMGDLLLLTGSLLDFK GYLGCQALSEMIQFYLEDVMPKAESDGEDIKEHVNSLGEKLTLLRRLRRCHQ FLPCENKSKAVEEVKSAFSLQERGVYKAMGEFDIFINYIEAYMTMKMRKN with polyhistidine tag at the C-terminus
Endotoxin level	<0.01 EU per 1 µg of the protein by the LAL method.
Activity	Measure by its ability to induce proliferation in MC/9-2 cells. The ED ₅₀ for this effect is <5 ng/mL.
Purity	>98% as determined by SDS-PAGE.
Form	Lyophilized
Storage Buffer	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile H ₂ O to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
Stability & Storage	This product is stable after storage at: <ul style="list-style-type: none"> • -20°C for 12 months in lyophilized state from date of receipt. • -20°C or -80°C for 1 month under sterile conditions after reconstitution. Avoid repeated freeze/thaw cycles.



SDS-PAGE analysis of recombinant swine IL-10

For research use only.