


PRODUCT INFORMATION

FGF-2 (Fibroblast growth factor, basic), Mouse

v. 231101

Catalog number	C02061-5UG / C02061-20UG / C02061-100UG
Package	5 μg / 20 μg / 100 μg
Description	FGF2, also known as a basic fibroblast growth factor (bFGF) and FGF- β , is a growth factor and signaling protein encoded by the FGF2 gene. FGF2 has been shown in preliminary animal studies to protect the heart from injury associated with a heart attack, reducing tissue death and promoting improved function after reperfusion. FGF-2 (bFGF) are also involved in a variety of biological processes, including embryonic development, morphogenesis, tissue repair, tumor growth, and invasion. Additionally, FGF-2 (bFGF) is frequently used for a critical component of cell culture medium, e.g., human embryonic stem cell culture medium, serum-free culture systems.
Source	Escherichia coli
Sequence	ALPEDGGAAFPPGHFKDPKRLYCKNGGFFLRIHPDGRVDGVREKSDPHVKLQL QAEERGVVSIKGVCANRYLAMKEDGRLLASKCVTEECFFFERLESNNYNTYRS RKYSSWYVALKRTGQYKLGSKTGPGQKAILFLPMSAKS with polyhistidine tag at the N-terminus
Endotoxin level	<0.1 EU per 1 µg of the protein by the LAL method.
Activity	Measure by its ability to induce 3T3 cells proliferation. The ED $_{50}$ for this effect is <1.5 ng/mL. The specific activity of recombinant mouse FGF-2 is approximately >1x 10^6 IU/mg.
Purity	>98% as determined by SDS-PAGE.
Form	Lyophilized
Storage Buffer	Lyophilized from a 0.2 μm filtered solution of PBS, pH 8.0.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile H_2O to a concentration not less than 200 $\mu g/mL$ and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
Stability & Storage	This product is stable after storage at: - 20°C for 12 months in lyophilized state from date of receipt. - 20°C or -80°C for 1 month under sterile conditions after reconstitution. Avoid repeated freeze/thaw cycles.





For Research Use Only.