

CXCL9 (C-X-C motif chemokine 9), Mouse

v. 231101

Catalog number	C02079-5UG / C02079-20UG / C02079-100UG
Package	5 µg / 20 µg / 100 µg
Description	<p>CXCL9, also named Monokine, is a member of the CXC chemokine family and is induced by gamma interferon (MIG). Following induced by IFN-gamma, this chemokine can attract T-cells. CXCL9 has close relationship with two other CXC chemokines named CXCL10 and CXCL11, additionally they all elicit their chemotactic functions by interacting with the chemokine receptor CXCR3. CXCL9 is also a cytokine that affects the growth, movement, or activation state of cells participating in immune and inflammatory response and work as a chemoattractant of activated T-cells.</p>
Source	<i>Escherichia coli</i>
Sequence	<p>TLVIRNARCSCISTSRGTIHYKSLKDLKQFAPSPNCNKTEIIATLKNQDQTCLDPD SANVKKLMKEWEKKINQKKKQKRGGKHKQKNMKNRKPPTQSRRRSRKTT with polyhistidine tag at the N-terminus</p>
Endotoxin level	<0.1 EU per 1 µg of the protein by the LAL method.
Activity	<p>Measure by its ability to chemoattract BaF3 cells transfected with mouse CXCR3. The ED₅₀ for this effect is <0.3 µg/mL.</p>
Purity	>98% as determined by SDS-PAGE.
Form	Lyophilized
Storage Buffer	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
Reconstitution	<p>It is recommended to reconstitute the lyophilized protein in sterile H₂O to a concentration not less than 200 µg/mL and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.</p>
Stability & Storage	<p>This product is stable after storage at:</p> <ul style="list-style-type: none"> -20°C for 12 months in lyophilized state from date of receipt. -20°C or -80°C for 1 month under sterile conditions after reconstitution. <p>Avoid repeated freeze/thaw cycles.</p>



SDS-PAGE analysis of recombinant mouse CXCL9

For Research Use Only.