

Catalog number	C15010-GMP-25KU / C15010-GMP-200KU		
Set package & Component	Cat.	Name	Amount
	C15010-GMP-25KU	GMP T7 RNA Polymerase (200 U/ μ L)	25 KU
		GMP 10X RNA Polymerase reaction buffer	1 mL
		GMP 100 mM DTT	1 mL
	C15010-GMP-200KU	GMP T7 RNA Polymerase (200 U/ μ L)	200 KU
		GMP 10X RNA Polymerase reaction buffer	4 mL
GMP 100 mM DTT		4 mL	
Description	Bacteriophage T7 RNA Polymerase is a DNA-dependent RNA polymerase with high specificity for the T7 promoter. This enzyme catalyzes the 5'→3' synthesis of RNA from DNA downstream from its promoter.		
Source	<i>Escherichia coli</i>		
Purity	>98% as determined by SDS-PAGE analysis.		
Unit Definition	One unit is defined as the amount of the enzyme incorporates 1 nmol of ATP into acid-insoluble product in 1 hour at 37°C.		
Endotoxin level	<0.05 EU per 1 μ g of the protein by the LAL method.		
Storage Buffer	T7 RNA Polymerase is supplied in 100 mM Tris-HCl (pH 7.9), 20 mM KCl, 1 mM DTT, 1 mM EDTA, 0.1% Triton® X-100 and 50% (v/v) glycerol.		
Storage	Store at -20°C for up to 6 months, and it is recommended to store at -80°C for long-term preservation. Avoid repeated freeze/thaw cycles.		
Reaction Condition	<p>1X RNA Polymerase Reaction Buffer, supplemented with 3 mM each ATP, UTP, GTP, CTP, and DNA template containing the T7 RNA Polymerase promoter. Incubate at 37°C.</p> <p>10X RNA Polymerase Reaction Buffer: 400 mM Tris-HCl (pH 8.0), 60 mM MgCl₂, and 20 mM spermidine.</p>		
Handling Instruction	For optimal storage, aliquot the enzyme, reaction buffer and DTT reagent into smaller quantities and store at recommended temperature. Avoid extended exposure to ice; instead, promptly retrieve the required portion and return it to the appropriate storage temperature.		
Manuel	<p>Standard RNA synthesis procedures:</p> <ol style="list-style-type: none"> 1. Below reaction mixture should be prepared under room temperature and combined in the following order: 		

Component	Amount	Final concentration
Nuclease-Free H ₂ O	X μ L	-
Template DNA	0.5-1 μ g	
GMP 10X RNA Polymerase Reaction Buffer	2 μ L	1X
ATP (100 mM)	0.6 μ L	3 mM
UTP (100 mM)	0.6 μ L	3 mM
CTP (100 mM)	0.6 μ L	3 mM
GTP (100 mM)	0.6 μ L	3 mM
GMP 100 mM DTT	2 μ L	10 mM
GMP T7 RNA Polymerase (200 U/ μ L)	1 μ L	-
RNase inhibitor (optional)	0.5 μ L	1 U/ μ L
Total reaction volume	20 μ L	-

2. Incubate at 37°C for 30 minutes to 2 hours.
3. Above reaction mixture may be scaled up or down proportionately.

Notes

1. Transcription reaction should be performed under RNase free condition. Use nuclease-free tubes, reagents, and water to avoid RNase contamination. Also, gloves when working with RNA.
2. To obtain optimal condition, NTP concentration can be titrated between 3 - 5 mM.
3. The volume of T7 RNA Polymerase can be titrated between 1-2 μ L in the IVT reaction to optimize your assay.

For Research Use Only.