


PRODUCT INFORMATION

Recombinant Mouse Activin A Protein, Tag Free

v. 240801

Catalog number	C02098-5UG / C02098-20UG / C02098-100UG
Package	5 μg / 20 μg / 100 μg
Description	Activin and Inhibin are members of the TGF-beta superfamily of cytokines and are involved in a wide range of biological processes including tissue morphogenesis and repair, fibrosis, inflammation, neural development, hematopoiesis, reproductive system function, and carcinogenesis. Activin is strongly expressed in wounded skin, and overexpression of activin in epidermis of transgenic mice improves wound healing and enhances scar formation. Activin also regulates the morphogenesis of branching organs such as the prostate, lung, and kidney. There is also evidence showed that lack of activin during development results in neural developmental defects.
Species of Origin	Mouse
Expression System	Escherichia coli
Sequence	Gly311-Ser426
Endotoxin level	<0.1 EU per 1 µg of the protein by the LAL method.
Activity	Measure by its ability to inhibit the proliferation of mouse MPC-11 cells. The ED_{50} for this effect is <10 ng/mL. The specific activity of recombinant human Activin A is approximately >1.0 x 10^3 IU/mg.
Purity	>95% as determined by SDS-PAGE.
Form	Lyophilized
Storage Buffer	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile H_2O to a concentration not less than 200 $\mu g/mL$ and incubate the stock solution for at least 20 min to ensure sufficient re-dissolved.
Stability & Storage	This product is stable after storage at: 20°C for 12 months in lyophilized state from date of receipt. 20°C or -80°C for 1 month under sterile conditions after reconstitution. Avoid repeated freeze/thaw cycles.





SDS-PAGE analysis of recombinant mouse Activin A, Tag Free

For research use only.